

## Expert View from Science

Anders Thapper, Stenbjörn Styring<sup>\*1</sup>, Guido Saracco, A. William Rutherford, Bruno Robert, Ann Magnuson, Wolfgang Lubitz, Antoni Llobet, Philipp Kurz, Alfred Holzwarth, Sebastian Fiechter, Huub de Groot, Sebastiano Campagna, Artur Braun, Hervé Bercegol<sup>\*2</sup> and Vincent Artero<sup>\*3</sup>

# Artificial Photosynthesis for Solar Fuels – an Evolving Research Field within AMPEA, a Joint Programme of the European Energy Research Alliance

**Abstract:** On the path to an energy transition away from fossil fuels to sustainable sources, the European Union is for the moment keeping pace with the objectives of the Strategic Energy Technology-Plan. For this trend to continue after 2020, scientific breakthroughs must be achieved. One main objective is to produce solar fuels from solar energy and water in direct processes to accomplish the efficient storage of solar energy in a chemical form. This is a grand scientific challenge. One important approach to achieve this goal is Artificial Photosynthesis. The European Energy Research Alliance has launched the Joint Programme “Advanced Materials & Processes for Energy Applications” (AMPEA) to foster the role of basic science in Future Emerging Technologies. European researchers in artificial photosynthesis recently met at an AMPEA organized workshop to define common research strategies and milestones for the future. Through this work artificial photosynthesis became the first energy research sub-field to be organised into what is designated “an Application” within AMPEA. The ambition is to drive and accelerate solar fuels research into a powerful European field – in a shorter time and with a broader scope than possible for individual or national initiatives. Within AMPEA the Application Artificial Photosynthesis is inclusive and intended to bring together all European scientists in relevant fields. The goal is to set up a thorough and systematic programme of directed research, which by 2020 will have advanced to a point where commercially viable artificial photosynthetic devices will be under development in partnership with industry.

**Keywords:** solar fuels, European Energy Roadmap, Joint Programming

**PACS® (2010).** 88.05.Ec, 82.47.Jk, 82.47.-a

**\*Corresponding author:** Stenbjörn Styring: Department of Chemistry-Ångström Laboratory, Uppsala University, Box 523; SE-751 20 Uppsala, Sweden. E-mail: stenbjorn.styring@kemi.uu.se

**\*Corresponding author:** Hervé Bercegol: CEA/DSM, Alternative Energies and Atomic Energy Commission, CEA Saclay, 91191 Gif-sur-Yvette, France. E-mail: herve.bercegol@cea.fr

**\*Corresponding author:** Vincent Artero: Laboratoire de Chimie et Biologie des Métaux, CNRS – UMR 5249, Université Joseph Fourier, CEA, 38000 Grenoble, France. E-mail: vincent.artero@cea.fr

**Anders Thapper, Ann Magnuson:** Ångström Laboratory, Department of Chemistry, Uppsala University, Uppsala, Sweden

**Guido Saracco:** Department of Applied Science and Technology, Politecnico di Torino, Torino, Italy

**A. William Rutherford:** Molecular Biosciences, Imperial College, London, UK

**Bruno Robert:** Institute of Biology and Technology of Saclay, CEA, CNRS and Université Paris Sud, Gif sur Yvette, France

**Wolfgang Lubitz, Alfred Holzwarth:** Max Planck Institute for Chemical Energy Conversion, Mülheim/Ruhr, Germany

**Antoni Llobet:** Institute of Chemical Research of Catalonia (ICIQ), Campus Universitari de Tarragona, Tarragona, Spain

**Philipp Kurz:** Institute for Inorganic and Analytical Chemistry, Albert-Ludwigs-University Freiburg, Freiburg, Germany

**Sebastian Fiechter:** Helmholtz Zentrum Berlin für Materialien und Energie GmbH, Institute for Solar Fuels, Berlin, Germany

**Huub de Groot:** Leiden Institute of Chemistry, Gorlaeus Laboratories, Leiden University, the Netherlands

**Sebastiano Campagna:** Department of Chemical Sciences, University of Messina, Messina, Italy

**Artur Braun:** Laboratory for High Performance Ceramics; Empa. Swiss Federal Laboratories for Materials Science and Technology; Dübendorf, Switzerland

This paper was issued from a workshop organised under the auspices of EERA. It should be considered as a collective work involving a much larger number of scientists than the named authors.

# 1 Background

## 1.1 Addressing social goals in energy research


Europe and most of its individual countries depend heavily on outside fossil energy resources. These are dwindling and their supply can easily be jeopardized. In addition, the use of the presently dominating energy carriers based on fossil fuels is connected to environmental and social factors that clearly affect the quality of life. In particular this concerns coal, oil and gas with severe availability constraints, pollution problems and alarming CO<sub>2</sub>-emissions. Catastrophes, in particular the damage done by the tsunami in 2011 in Japan, have increased the public worries about large future nuclear power programmes, and in addition significantly increased the projected costs of new nuclear power plants.

## 1.2 Organisation of European energy research

These factors have prompted the European Union to accept that an energy transition will occur and enter into

a planning phase the aim of which is the progressive replacement of fossil fuels by clean energy, mainly from renewable resources. This long term vision has two time-scales. In the near and mid-term future, the 27 member states have already set targets for the share of renewable energy in the overall consumption in 2020, with an average of 20% for the entire Union. Enacted as an EU directive [1], this goal is part of the Strategic Energy Technology Plan [SET-Plan]. For the longer term, an energy roadmap has been drawn up all the way to the year 2050 [2]. It calls for a reduction of more than 80% in greenhouse gas emissions, for a strong commitment to enhance energy efficiency and for efforts to increase the penetration of renewable energy to reach all sectors of society. The use of renewable energy is intended to reach levels higher than 50% of the total European energy consumption by 2050.

Today the 20% renewable energy target for 2020 appears reachable, provided that the installation of facilities producing renewable energy continues to increase at the current rate. From 2004 to 2010, the share of renewable energy in Europe experienced an annual growth of about 8% (Figure 1). If this growth continues at the same rate, it has the potential to bring Europe above the target set for 2020 (although this certainly requires substantial



**Fig. 1:** Share of renewables in primary energy consumption in the European Union (in %) plotted on a log scale – data from Eurostat and from the Swiss federal office of energy for the period 2004–2010. In most European countries (we show here only the statistics for the home countries of the authors of this contribution), renewable energy was promoted from the beginning of the century leading to a steady development. The present pace permits some of the targets defined for 2020 to be reached [1]. After 2020, scientific and technological breakthroughs are needed, especially if renewable energy use is to continue increasing to levels higher than 50% for the whole EU in 2050. For this, the production of solar fuels by Artificial Photosynthesis would represent a key breakthrough. It is time now to invest in an Artificial Photosynthesis roadmap, if we want to maintain or even accelerate the present trend in the energy transition.

financial investments). However, in order to maintain the pace of the energy transition until 2020 and beyond, there is a real need for a major research effort in many energy sectors. These have to address basic and applied topics concerning both the organisation and the technology of the entire energy sector.


Together with the launch of the SET-Plan in 2008, European public research and technology organisations were encouraged to unite their scientific competences: ten founding partners constituted the European Energy Research Alliance (EERA) [3] and started to work at defining and implementing Joint Programmes designed to accelerate research towards low carbon technologies and to rationalise investments. The alliance now includes more than 150 participating institutions throughout Europe. Its *raison d'être* is threefold: a) to minimize redundancy and duplication in research; b) to concentrate funding on the best performing teams; c) to build trans-national teams of a critical size.

To date, EERA has launched 13 Joint Programmes. Most of them are focussed on wind, solar (thermal and photovoltaic), biomass, geothermal, and ocean energy and the technologies crucial for their implementation (for example smart grids). In particular the Joint Programme "Energy Storage" aims at improving and tuning efficient large-scale storage capacities for grid-connected wind or solar facilities. Chemical storage, for example the conversion and storage of solar energy in the chemical bonds of hydrogen ( $H_2$ ) made from water, is being increasingly implemented in the so-called "Power-to-Gas" scheme. Considering the 2050 road-map [2], it will with time become necessary to convert an increasingly larger share of the renewable energy into gaseous or liquid fuels. One important reason is that most of the energy consumption in Europe (and globally) is not in the form of electricity, which is the product of wind-power, solar cells, concentrated solar heat and several other renewable technologies. Instead, more than 80% of the primary energy is used mainly as fuels to drive a vast variety of processes in our societies (see e.g. Eurostat) [4]. A second reason is limitations of the electricity grid, which prevents very large penetration of renewable electricity coming from intermittent energy sources often located far away from the final users. One promising solution to both these problems is Artificial Photosynthesis which provides all of the energy supply, the necessary energy storage and the facilitated transportation. In addition, Artificial Photosynthesis allows utilisation of renewable energy at both the local and the continental level producing fuels made from solar energy and water.

### 1.3 Solar fuels and artificial photosynthesis

The introduction of CO<sub>2</sub>-lean, transportable energy carriers made from renewable resources would directly minimize the use of fossil fuels. Unfortunately, biomass-based fuels (although heavily developed and dominating today) can only be important on a restricted scale. Globally there will not be enough biomass to allow for biofuel production to exchange for the fossil fuels. This holds today and even more so in the future when the global energy demand has increased further. Solar energy represents by far the largest of the presently useful renewable energy flows. Therefore, the vision of *solar fuels as important future energy carriers* makes this a central research theme that is quickly moving up the global agenda. The terms "solar fuel", "artificial photosynthesis," and "artificial leaf" have become established in recent years. All of them reflect the same goal, to make a fuel from solar energy and sustainable raw materials, in particular using water as the electron source (Figure 2).

In the long run, new scientific breakthroughs will be needed for solar fuels or other future emerging renewable energy technologies to become used on a scale that is necessary to replace fossil fuels. Basic, multidisciplinary science is required in order to push forward the energy transition by the second half of the century. In 2008, the European Science Foundation published a science policy briefing calling for an action plan towards the development of solar fuels [5]. The ESF later sponsored a EUROCORES programme for solar fuels, which is currently funding two solar fuel networks dealing with some aspects of the scientific road-map to solar fuels [6]. To increase the impact of innovation coming from academic research, EERA recently started a basic science initiative: the Joint




**Fig. 2:** Overview of artificial photosynthesis. The vision is to produce a solar fuel from solar energy and water. There are different solar fuels but they are always made with solar energy as the only energy source and water as the electron source. The science in AMPEA Artificial Photosynthesis is intended to develop direct energy conversion processes where water is the raw material. The box represents the many challenging research subfields involved. From an applied perspective the box represents the future systems or devices, which will be producing the solar fuel from solar energy and water.

Programme AMPEA, standing for Advanced Materials & Processes for Energy Applications. AMPEA was launched at the end of 2011 and to date more than 40 European scientific institutions participate in the programme. In Parts 2 and 3 of this paper, we will describe how the application of the sub-field “Artificial Photosynthesis” will be developed within AMPEA.

In recent years there have been calls for investments in science in order to obtain breakthroughs on energy technologies. Steven Chu, Nobel Prize laureate in Physics and present Secretary of Energy of the United States, said that on the subject of energy that he was “optimistic that science can offer better solutions than we can imagine today . . . if we harness the creativity and . . . intellectual horsepower of our best scientists in the right way” [7]. The Department of Energy (DOE) organised a consultation of national and international experts which produced orientation documents and eventually gave birth to new funding schemes for energy science. DOE funded a series of Energy Frontier Research Centers of which several have large solar fuels activities. DOE also created the so-called Energy Innovation Hubs in 2010, and among them, a Joint Centre for Artificial Photosynthesis (JCAP) [8] was established between the California Institute of Technology and the Lawrence Berkeley National Laboratory in California. Currently this is probably the world’s largest operation (two dedicated laboratory buildings, one at each campus) on artificial photosynthesis purely aiming at the development of solar fuels. In Europe, the Swedish Consortium for Artificial Photosynthesis (CAP) was the first large collaboration of this kind [4]. The Swedish CAP started much earlier, in 1994, and is since 2005 concentrated at Uppsala University which hosts the largest laboratory entirely devoted to solar fuels research in Europe. In Europe several new efforts of this dedicated type are being launched in for example Italy [9] and The Netherlands with the BioSolar Cells consortium [10]. Also elsewhere in the world, especially in Japan, Korea, China and Australia, ambitious programmes have been initiated. This burgeoning activity reflects that solar fuels is one of very few options that mankind can develop to achieve and maintain a sustainable future for the 9–10 billion people anticipated to be living on the planet by the year 2050 [5].

#### 1.4 AMPEA and artificial photosynthesis

AMPEA has a matrix structure which is shown in Figure 3. The science, intended to drive the European energy research in areas where the basic science content is high, will be carried out in energy applications. The applica-



**Fig. 3:** The structure of AMPEA is a matrix of tools and applications aimed at fostering the development of future emerging energy technologies, while favouring the use of scientific tools provided by researchers of the Applications domains. AMPEA Artificial Photosynthesis is described in the present text. Other emerging applications are currently being defined.

tions are crossing three technical sub-programmes (SP) that involve basic scientific tools. For SP1 these are the design and synthesis of new materials; for SP2 the physical modelling of materials, devices and processes; and for SP3 the characterization of materials, devices and processes. The technical sub-programmes constitute a core base and are intended to deepen the research carried out in the scientific applications. Artificial Photosynthesis is the first of these applications. Its creation (Figure 3) within the AMPEA Joint Programme is intended to boost research in this area on a pan-European basis. Its immediate inclusion in the first phase of the AMPEA Joint Programme reflects its urgent character. Artificial Photosynthesis in AMPEA should be inclusive and gather all scientists active in related fields in Europe. This research community is diverse and already quite large. It is rapidly growing by attracting young researchers and the field in Europe is estimated to already involve as many as 100 internationally recognized PI’s in more than 15 countries.

*A solar fuel is always made using solar energy as the only energy source.* Huge amounts of solar energy are available everywhere and are often abundant, even in densely populated areas. In contrast to other renewable energy options there is more than enough solar energy available to allow the shift from fossil fuels, even at fairly modest conversions efficiencies of the solar energy to the desired product (a solar fuel or solar electricity). Even in

Northern Europe there is enough solar energy to contribute substantially to the energy system in the summer with the long days. However, the solar energy is lacking in winter when energy demand is high. Therefore, solar energy can become a truly important energy source also in these very northern countries only through its conversion into a storable energy carrier, the solar fuel.

For the widespread and cost effective development of solar energy in Europe, it is critical that very efficient processes are developed. In particular, it is urgent to develop high efficiency artificial photosynthesis routes for the *direct conversion* of solar energy into fuel [4, 9]. Such routes are not yet available and the AMPEA Artificial Photosynthesis Application is created specifically to address this point. Indirect processes for fuel production, using for example electricity or biomass as intermediary energy carriers, appear – even though they work – less optimal because of inherent energy losses. They are not part of AMPEA but are studied partially by other Joint Programmes of EERA. Synergies shall be profitably developed between these parallel paths towards solar fuels, on catalysts, electrolytes, materials synthesis, design of devices etc.

*The raw material for the fuel* is the second, equally important, key aspect of this process. Artificial photosynthesis targets *water as the raw material* [4]. Thus, AMPEA Artificial Photosynthesis gathers basic and applied science with the aim of providing a CO<sub>2</sub>-lean fuel based on solar energy and water (Figure 2). We envision extensive development of artificial photosynthesis for solar fuels as a replacement for fossil fuels thereby avoiding the effects on the climate caused by the current extensive use of fossil fuels. The road to success will be long and winding, the stakes are high but the potential benefits enormous. Scientific breakthroughs in this area could quickly be transformed into technology, lead to new industrial sectors, and drive positive societal changes. At this moment the field of solar fuels in Europe is scattered. AMPEA Artificial Photosynthesis is designed to change this.

*The intended solar fuel should address the market demand that could vary in different countries or regions.* Presently there are only a few ideas for solar fuels that are being pursued internationally. AMPEA Artificial Photosynthesis intends to gather the available science in Europe to design several promising routes for fuel production from solar energy and water [4]. Many scientists target hydrogen as the solar fuel. This is driven by intentions in for example the European road map towards the hydrogen society (in which hydrogen should be non-carbon based by 2050) and is a natural choice when water is the raw material. Proof of concept principles already exists and it is possible to produce hydrogen, even at high pressure,

with artificial photosynthesis. However, other solar fuels, such as carbon containing solar fuels like alcohols produced from concentrated CO<sub>2</sub> sources, are also potential targets [9, 11].

A key element in the research is *solar energy conversion*. AMPEA Artificial Photosynthesis aims to use light energy to drive all reactions leading to fuel formation and to develop artificial photosynthesis in supra-molecular systems and in photo-electrochemically active nanostructured materials. Molecular sensitizers or semiconductor materials will be employed to harvest the light and to convert the energy into an electrochemical potential able to drive electrons between catalysts. All components have to be made from cheap and abundant elements and the system must drive both the efficient oxidation of water and reductive processes (hydrogen evolution and/or reduction of carbon dioxide) to generate the fuel. A core activity to achieve this is the design and synthesis of stable, cheap and efficient catalysts for water oxidation and fuel production. A central feature of such catalysts is their ability to perform advanced electron and proton management. Advances in this field will benefit from suppression by detailed biochemical, structural and biophysical knowledge of enzymes that carry out these processes in nature (hydrogenases, water oxidizing enzyme and photosynthetic reaction centres) and biology will also serve as inspiration in this field. The integration of the knowledge from functional natural systems into man-made catalysts is a key aspect of this programme.

*Energy relevance – strategic objective.* The strategic objective of AMPEA Artificial Photosynthesis is to meet demands for an exchange for fossil fuels by the production of solar fuels from solar energy using water as the electron source. When this is accomplished, solar energy will be converted to chemical energy in a fuel. The fuel can be stored, from summer when solar energy is abundant to winter when demand is high, from daytime with strong sunshine to evenings and nights etc. It can be transported to where it is needed. It has the potential to meet the present need of ca. 80% fuel in the energy consumption. This research is complex, challenging and without shortcuts. It is consequently long-term and reaches into a time-domain beyond 2020. It demands work in cross-disciplinary constellations of researchers where expertise from many different fields join forces. AMPEA Artificial Photosynthesis addresses an area where Europe has an edge with respect to other international research efforts. The programme will create new interfaces where strong synergies can be anticipated. Breakthrough results and unexpected quantum leaps in the science are expected to be promoted at these interfaces allowing development

towards large-scale application through the new collaborations explored in AMPEA. The programme integrates a broad arsenal of scientific techniques and thinking/knowledge from scientists with many competences. The planned research is fundamental but with a strong emphasis on addressing the total conversion chain, by integration of parts and modules to establish functional devices, first on the laboratory scale and later on the industrial scale.

AMPEA Materials sub-programme SP1 aims at rationalising routes of materials synthesis, and materials assembly in the framework of nanotechnologies: for Artificial Photosynthesis, benefits will concern the concept of electrodes and membranes, the integration of catalysts properties and the optimisation of interfaces that are main sources of energy loss. An important feature in AMPEA is the merging of materials science with the study of physical and chemical processes within the modelling and characterisation sub-programmes SP2 and SP3. The complex multidimensional problems of energy solar fuel producing devices will be investigated in specific platforms, by developing specific modelling tools and by using dedicated large-scale instruments. In situ operando characterisation will give insight into the materials and processes at the heart of the photo-catalytic devices designed for artificial photosynthesis.

rial. While these biofuels might be very useful in certain regions or countries, they are always limited by the availability of the corresponding raw material. This limits their application on a global scale and such methods are *not covered* in this programme. Instead, the intended raw material must be essentially inexhaustible, cheap and widely available [4]. There are few options and *the target in this proposal is to develop processes where water is the raw material*. The question of the availability and cost of clean water can be raised. However the water usage in processes for solar fuels production using artificial photosynthesis is orders of magnitude smaller than for agricultural processes [12]. The amount of water used in the process is therefore of little concern from an environmental perspective. Processes where water is split into its constituents by solar energy can become large contributors to shift away from fossil fuels on a global scale. The development of technologies for direct solar to fuel conversion requires revolutionary breakthroughs in several areas of fundamental science. Equally important is the integration of the different parts and modules in smart (responsive) matrices at an early stage and to produce prototypes and functional devices. An additional mind-provoking possibility is to use semi-artificial components from natural species that are engineered for superior performance beyond what nature has to offer.

## 2 Objectives

### 2.1 Long-term impact: a conceptual change of today's fossil based energy system on the terawatt scale

#### 2.1.1 Energy source and raw material


Solar fuels are made using *solar energy as the sole energy source and water as the raw material*. They will consequently be able to provide CO<sub>2</sub>-lean, renewable and storable energy carriers in almost unlimited quantities. Today, functional and cost-efficient direct technologies for solar fuels production are not available and the necessary science to change this is the focus of AMPEA Artificial Photosynthesis. The long-term vision is that the development and use of solar fuels shall become a major factor to alleviate the effects of the extensive use of fossil fuels, thereby providing a paradigm shift away from our present fossil-based energy system.

For 1<sup>st</sup> or 2<sup>nd</sup> generation biofuels (the presently dominating renewable fuels), biomass or waste is the raw mate-

### 2.2 Scientific objectives and linked research areas

*AMPEA Artificial Photosynthesis will focus on direct processes for solar fuel production involving several research subfields.* It will provide great possibilities for breakthrough research and unexpected, rapid progress. Novel synergies will emerge between the research in the scientific applications and scientists in the technical subprogrammes of the AMPEA Joint Programme.

Figure 4 shows an overview of the science and sub-fields in AMPEA Artificial Photosynthesis and Figure 5 illustrates a few representative scientific details from the respective subfields. For a long time, the most critical hurdle to be overcome has been the design of a stable, light driven catalyst that can oxidise water at sufficient rates. The catalysts for water-oxidation and fuel production can be *molecular and/or solid state or nanomaterials*. They share similar physical principles and similar, major scientific challenges are encountered in multi-electron catalysis and proton-coupled electron transfer. For both molecular and nanoscaled catalysts, water-oxidation is a major challenge. To make the huge quantities of solar




**Fig. 4:** Overview of the broad scope of the three subfields in AMPEA Artificial Photosynthesis. Numerous breakthroughs probably lie in the collaboration between the three subfields, in the design of hybrid devices as well as in the understanding of basic principles of water oxidation, hydrogen production, light absorption etc. Knowledge transfer from biological systems that function extremely efficiently is a key element in AMPEA Artificial Photosynthesis. The development of the necessary breakthrough concepts will be facilitated by the connection of Artificial Photosynthesis to the technical sub-programmes that bring a wide range of scientific methodologies; abilities to optimize design and synthesis etc. (see e.g. Figure 3).

fuels needed for a transition from fossil fuels it is necessary that the catalysts are made from earth abundant elements like manganese (in natural photosynthesis) [13], iron [14] or cobalt [15] while the use of scarce and expensive elements such as ruthenium, platinum or iridium otherwise might become a prohibitive bottleneck [16]. There are few efficient molecular catalysts based on these

abundant elements and development of such catalysts is therefore in the early focus for many research efforts of the AMPEA participants. However, there exist heterogeneous catalysts based on cobalt [15, 17, 18] manganese [18, 19] or iron that are ready to try in first-generation device-oriented research. Such attempts will allow fundamental, in-depth understanding of physical and chemical processes in working artificial photosynthesis devices. The same arguments also hold for available efficient molecular catalysts based on ruthenium which are useful model systems both for basic and applied research [20–22]. To consolidate our basic knowledge and to accelerate the research, we will not wait until the last generation of catalysts are developed. In all concepts, the fuel will be produced utilizing the reducing equivalents made available from the light-driven oxidation of water. The fuel can be hydrogen obtained by the reduction of protons, but also the reduction of carbon dioxide ( $\text{CO}_2$ ) can yield fuels such as alcohols. The latter might be easier to use with current technology but is probably more difficult to make in a direct process (although for instance methanol can be efficiently made from hydrogen in industrial processes).

New frontier science is created in AMPEA Artificial Photosynthesis by linking the *three subfields* overviewed in Figures 4 and 5:

- *A. molecularly designed systems*
- *B. solid-state components*
- *C. nature-guided design*



**Fig. 5:** Overview of AMPEA Artificial Photosynthesis high-lighting the cross-disciplinary nature of the research. Upper part: the solar fuel forming subfields. Right: artificial photosynthesis in devices that make a fuel from solar energy and water using light-absorbers (S) linked to light-driven molecular catalysts made from Mn, Co, Ni or Fe. Left: man-made nanostructured photo-electrochemical systems, here symbolized by hematite (iron-oxide) and a Mn/Ca layer structure used in promising modular assemblies. Lower half: The science stands on solid ground and is built on studies of photosynthetic reaction centres and light harvesting complexes (middle – Photosystem II with antenna and water oxidizing complex), hydrogen metabolizing enzymes (right Ni-Fe hydrogenase). Strong (bio)physical knowledge of solar energy conversion is a key element to the programme. A catalytic system for proton reduction based on carbon nanostructures with a linked catalyst is also shown. Shaded triangles symbolize overlapping areas created in AMPEA Artificial Photosynthesis by synergies with the technical sub-programmes. Some of these will be subject for projects in future calls.

All have strong synergies with the technical sub-programmes (SP 1–3, Figure 3) thereby making AMPEA Artificial Photosynthesis an ideal starting point for immediate activities in application-oriented basic research on renewable energy. In particular, the joint efforts of the field will strive to demonstrate the first generation of viable devices for technical application within the time-frame of the Horizon 2020 framework programme [23].

### 2.2.1 Sub-field A

The first subfield concerns the use of *molecularly designed systems* (Figures 4 and 5, purple, right). The catalysts are often developed following bio-inspired approaches, but might also contain abundant elements not used by biological systems [24]. The four-electron oxidation of water is clearly one of the main research challenges and a bottleneck for the successful development of artificial photosynthesis. For water oxidation, manganese containing catalysts [25–27] mimicking key aspects of the water oxidizing chemistry in the photosynthetic enzyme Photosystem II [13] and recently introduced cobalt catalysts [15, 17, 28, 29] are promising tracks to follow. There exist also efficient ruthenium catalysts for water oxidation. Although based on a rare element, these are useful to study to increase our knowledge about possible water oxidizing mechanisms and also for early device-oriented work. Two-electron molecular photo-catalysts for hydrogen formation will be based on abundant, cheap transition metals (instead of rare noble metals like platinum) [30]. Biomimetic di-iron and nickel-iron systems mimicking the active site in hydrogenase enzymes [14] are promising but, there are also very useful cobalt-based catalysts [15, 31]. Additionally, the development of stable dye systems for light-harvesting and sensitization are important issues which need attention and evaluation [20, 30]. Within the time-frame for Horizon 2020, the research in this subfield will develop second-generation devices for technological evaluation.

A strong advantage with molecular catalysts is that it is easier to achieve very high degrees of mechanistic understanding than in other conceivable systems. The molecular catalysts can be rationally tailored, tuned and studied for mechanism and performance using a multitude of sophisticated spectroscopic techniques, many available within the technical sub-programmes of AMPEA. This is necessary both for an incorporation of molecular catalysts and dyes into functional devices and also for the design and development of next-generation components.

### 2.2.2 Sub-field B

For the developments of *solid-state components* (second subfield; Figures 4 and 5, red left) AMPEA Artificial Photosynthesis will investigate solar-driven hydrogen production via photo-electrochemical water splitting [32, 33]. Thin-film photovoltaic materials can be envisioned as short or medium-term solutions if corrosion issues can be solved through the development of protective mechanisms or passivation layers [34]. However, the long-term target should be to construct a cheap nanostructured semiconductor device based on earth-abundant elements [35]. Novel design strategies for the coupling of inorganic components to catalysts need nanotechnology and material science to prepare innovative composite structures in which each component performs its specialized function. Again, one of the key issues in this context is the design of *catalytic nano-particles* or incorporation of *molecular catalysts* that will be interfaced with semiconductors to catalyse water oxidation and hydrogen (fuel) formation. High surface areas to maximise the contact between the catalyst and the substrate and low overvoltages for the desired reactions are required. A critical factor here is the interaction with the technical sub-programmes within AMPEA regarding the scalable production of catalysts and structured components. This also holds for establishment of new methods for elucidating elementary steps of catalysis on solid-state surfaces.

A promising concept is the development of mixed systems, for example by combining solid-state scaffolds with molecular light-driven catalysts. These should preferentially be based on abundant materials like iron, manganese, nickel or cobalt. This development will be initiated and facilitated by the close interaction of the researchers within AMPEA Artificial Photosynthesis.

### 2.2.3 Sub-field C

The third subfield involves *nature-guided design* (Figure 4 and 5, green middle) where we learn from Nature in our attempts to make entirely man-made chemical systems. Knowledge transfer from efficient photosynthetic systems and enzymes suppress and will strongly fertilize AMPEA Artificial Photosynthesis. This subfield also includes several aspects concerning *semi-synthetic/hybrid systems* involving both biological and man-made components [36, 37]. AMPEA Artificial Photosynthesis invites also this science to be part of the initiative. Importantly, knowledge gained from biological principles and the nature-guided design of devices is envisaged as a driving force and inspi-

rational source also for the research teams with no biological background. The understanding and learning from the chemistry of the natural oxygen evolving complex [28] and hydrogenases [38], for example concerning issues with electron and proton management at the catalytic centers [39–41], is of key importance. These processes are poorly understood so far in the context of materials for artificial photosynthesis. The very high quantum yield observed in natural systems may involve proteins as smart (responsive) matrices which reflect the tight coupling between the chemical reaction and the protein motions [42]. The transfer of principles from biological systems to artificial materials and devices will be an important task of the subfield nature-guided design.

### 2.3 Devices, demonstrators, prototypes — beyond the state of the art

Many necessary chemical building blocks like catalysts and charge separating ensembles have become available in recent years. Several are made and described by future participants in AMPEA Artificial Photosynthesis. The great challenge in AMPEA Artificial Photosynthesis, and its central research goal, is the assembly and optimization of the different parts to construct half-cells and thereafter functional integrated modular systems to assemble complete photo-catalytic devices that can operate with good yield against wasteful back reactions. Few systems like this have been constructed so far [9, 43] but success here would be ground-breaking for solar fuels production. This core element is virtually lacking from earlier research on artificial photosynthesis.

The idea in AMPEA Artificial Photosynthesis is to bring together many existing pieces from various European groups and laboratories and combine these in different device concepts. The aim is to directly connect photosensitizers to catalysts and surfaces or catalysts with semi-conductor materials and to study the half-cells constructed in this way using theoretical, chemical, physical methods as well as implementing advanced engineering knowledge for the first time in this part of the field.

Since there are many different components to choose from (many catalysts, different surfaces and matrices, many photosensitizers) many concepts will be possible and can be explored in such a programme on a European scale. This is the target of the programme and the intention is to have a significant number of laboratory scale pilot devices in operation just a couple of years from now. At this stage it is not useful to make bold promises or define exact goals for efficiencies, durability, costs etc.

since the basic science content in the work is by necessity high. However, all work must be pursued with such targets in mind and a fruitful development in the field would be that such targets could be reliably introduced towards the end of the Horizon 2020 programme period.

### 2.4 Milestones

The core milestones for the AMPEA Artificial Photosynthesis are: i) To build up a joint continental network in artificial photosynthesis research, involving a wide range of sciences with a common goal. ii) To accelerate and focus the European science in this challenging field with global impact more efficiently than can be achieved by national or individual efforts. iii) To prove the principle in a series of device concepts on a time scale of a few years. At this stage, all sub-fields are equally important and each sub-field (Figures 4 and 5) has internal, traditional scientific milestones.

### 2.5 Synergy between programme elements

The science in AMPEA Artificial Photosynthesis has a broad scope. All methods depend on advanced chemistry at modest temperatures and mild conditions. Larger physical scales than molecular interactions are also crucial in the control and long term durability of the devices. It is anticipated that much progress will be achieved at the new interfaces created between the scientists in AMPEA Artificial Photosynthesis working together and creating new research interactions with scientists in the technical sub-programmes of AMPEA. Multiscale integration and multifaceted understanding are the main goals of the technical sub-programmes. The central goal for Artificial Photosynthesis is the development of functional half-cells or entire solar-fuel cells where for example molecular catalysts are merged with semiconductor surfaces or photo-active material surfaces. Studies of this kind of devices demand interactions between scientists that seldom have collaborated but have complementary expertise. A required target for the entire scientific field of artificial photosynthesis is to address the challenge to provide renewable energy on the terawatt (TW) scale. This will only become possible to reach if the new synergies created within AMPEA Artificial Photosynthesis can be developed and fully exploited. This involves joint workshops between subfields where synergies are emerging or already strong. A useful development would be the creation of a European conference line. This could be timed to years when the Gordon research conference on solar fuels is not held.

### 3 Roadmap for an artificial photosynthesis network

Artificial photosynthesis has already been the subject of a few seminal European networks (e.g. Ru-Mn for artificial photosynthesis, SOLAR-H, SOLAR-H<sub>2</sub>, Solhydromics, nanoPEC, EuroSolarFuels, ArtipHyction, PERSPECT-H<sub>2</sub>O, H<sub>2</sub>-NanoSolar etc.) and initiatives in several countries provide a fertile ground for further development. To strengthen this important research field on the European level, knowledge from all European countries was brought together on the occasion of a workshop in Mülheim/Ruhr in Germany (October 2012) to achieve the long-term goal of mastering artificial photosynthesis. Their combined expertise at this workshop covered so diverse fields as biochemistry, molecular and supramolecular chemistry, photo-catalysis, photoelectrocatalysis, electrochemistry, material science, theoretical chemistry, optoelectronics, engineering etc. The AMPEA workshop gathered most of the important players of this rapidly developing scientific field on a European level.

The outcome of the meeting was: 1) the identification of bottlenecks and the formulation of milestones for artificial photosynthesis; 2) the definition of the scientific content and vision of AMPEA Artificial Photosynthesis; 3) the outline of a European road map to overcome the scientific and technological hurdles on the way to efficient solar energy conversion into solar fuels on a very large scale; 4) the definition of the necessity to fund this road map on a continental scale to a level required to be competitive with related initiatives outside Europe; 5) the foundation of a European Solar Fuels Society intended to be an organisational platform for the field in Europe.

The scientists participating in the workshop discussed the science within different time frames with respect to the time defined by the SET-plan and where the field currently exists in Europe. This is an on-going discussion and the points described below only present a starting point for the creation of a more detailed and broader road map for the development of solar fuels. The next AMPEA plenary meeting in this process is scheduled for June 11–12, 2013 in the Netherlands.

#### 3.1 Short timescale (2013/2014)

There are no quick fixes in solar fuels research and most of the research is long term. Despite this, an important issue on the short time frame is to organize the field in Europe in an inclusive and scientifically broad manner. It is imperative that both large scientific organisations and smaller

groupings in single universities or from smaller countries are welcome and can be accommodated in the networks and programmes created. The specific target is the scientific integration of the technical sub-programmes and the Artificial Photosynthesis Application, which began at the Mülheim workshop and shall be furthered in a joint AMPEA workshop in spring 2013. An important outcome will be the creation of the first calls for proposals in Artificial Photosynthesis within Horizon 2020, the next framework programme for EU research funding. It is also necessary to establish with relevant EU representatives the importance of Artificial Photosynthesis for the EU initiative leading to the “third industrial revolution” [44].

Early discernible bottlenecks and thereby core agendas in the research programmes include: improving both molecular and solid state catalyst efficiencies and stabilities, mechanistic understanding of critical components, controlling electron and proton management, efficient light absorption and charge-separation. The science involved here is both fundamental and application oriented.

#### 3.2 Medium timescale (2015 to 2020)

During this time-frame a programme, jointly funded by member states and the European Union, will develop a series of Artificial Photosynthesis devices on a laboratory scale based on our various technologies. These first generation devices will help to identify new and still remaining challenges and pin-point factors to focus future efforts on. These devices shall take advantage of the regular improvement of molecular and materials design, shall initiate new multiscale modelling actions and get retrofit from them, and shall be the subject of thorough investigations with dedicated instruments and lines of large scale facilities.

During this period it will also be important to develop a second target involving integration of molecular catalysts and solid state matrices in devices, as well as to implement the use of nature-guided approaches. This science will cover mechanistic studies, nano-structuring, light management, interface engineering, smart (responsive) matrices, grafting and assembling of modules and is well suited for efforts within the AMPEA Joint Programme.

Other valuable areas to be developed have to deal with standardisation and agreement on how to compare device performances as well as taking initiatives to develop a common scientific terminology for the field through summer schools, conferences and the writing of joint texts.

The most important goals and commitments for 2020 shall be the availability at this date of methods and devices

that industry can begin to consider for a commercial deployment.

### 3.3 Long-term (2020 and beyond)

At this time the field will have matured and there will be more scientists and approaches involved. The research under Horizon 2020 will enable a strong interaction between academic science and industry and thus allow the development of technologically more advanced devices, testing and evaluation towards commercialisation and life-cycle analyses. It will be possible to approach the 10%-efficiency goal for the conversion of solar energy to fuel. The real target is the development of efficient, long-lasting and cost effective device concepts for solar fuels such that direct conversion of sunlight into CO<sub>2</sub>-lean fuels is an accepted and viable part of a future energy mix in Europe. Parallel to this the basic science is now coupled to strong efforts in application oriented research and development driving the first generation of real solar fuel devices towards up-scaling for commercial competitiveness.

Received: January 21, 2013. Accepted February 4, 2013.

## References

- [1] Directive 2009/28/EC of the European Parliament and of the Council of 23 April 2009 on the promotion of the use of energy from renewable sources. [cited Jan. 8, 2013]. Available from: <http://eur-lex.europa.eu/LexUriServ/LexUriServ.do?uri=OJ:L:2009:140:0016:0062:en:PDF>
- [2] Communication from the Commission to the European Parliament, the Council, the European Economic and Social Committee and the Committee of the Regions – Energy Roadmap 2050 /\* COM/2011/0885 final \*/. [cited Jan. 8, 2013]. Available from: <http://eur-lex.europa.eu/LexUriServ/LexUriServ.do?uri=CELEX:52011DC0885:EN:NOT>
- [3] The European Energy Research Alliance [homepage on the Internet, cited Jan. 8, 2013]. Available from: <http://www.eera-set.eu>
- [4] Styring S. Artificial photosynthesis for solar fuels. *Faraday Discuss.* 2012;155:357–76.
- [5] Harnessing Solar Energy for the Production of Clean Fuel, Science Policy Briefing 34 of the European Science Foundation, September 2008. [cited Jan. 8, 2013]. Available from <http://www.esf.org/>
- [6] Pandit A, Holzwarth A, de Groot H. editors. Harnessing Solar Energy for the Production of Clean Fuel, ESF White Paper, ISBN 9789090239071 (2008). [cited Jan. 8, 2013]. Available from <https://openaccess.leidenuniv.nl/handle/1887/13349>
- [7] Quoted from Steven Chu's Budget testimony on the energy hubs, June 3, 2009. Washington, DC: U.S. Department of Energy [cited Jan. 8, 2013]. Available from <http://www.ne.doe.gov/AdvModelingSimulation/energyinnovations.html>
- [8] California Institute of Technology, Joint Center for Artificial Photosynthesis (JCAP), [homepage on the Internet]. Berkeley, CA, USA [cited Jan. 8, 2013]. Available from <http://solarfuelshub.org/>
- [9] Bensaid S, Centi G, Garrone E, Perathoner S, Saracco G. Towards Artificial Leaves for Solar Hydrogen and Fuels from Carbon Dioxide. *Chemsuschem.* 2012;5(3):500–21.
- [10] <http://www.biosolarcells.nl/en/>
- [11] Morris AJ, Meyer GJ, Fujita E. Molecular Approaches to the Photocatalytic Reduction of Carbon Dioxide for Solar Fuels. *Acc Chem Res.* 2009;42(12):1983–94.
- [12] It can for example be calculated that making hydrogen from all the water carried by the water stream Ätran in Sweden would be sufficient to cover the entire energy need by the present population in Europe. It has also been calculated that the splitting of about one Olympic swimming pool per second would be sufficient to cover the entire energy need of the world in 2050 assuming a doubling of the current energy use ([http://poptech.org/popcasts/dan\\_nocera\\_personalized\\_energy](http://poptech.org/popcasts/dan_nocera_personalized_energy) [cited Jan. 8, 2013])
- [13] Umena Y, Kawakami K, Shen JR, Kamiya N. Crystal structure of oxygen-evolving photosystem II at a resolution of 1.9 angstrom. *Nature.* 2011;473(7345):55–U65.
- [14] Tard C, Pickett CJ. Structural and Functional Analogues of the Active Sites of the [Fe]-, [NiFe]-, and [FeFe]-Hydrogenases. *Chem Rev.* 2009;109(6):2245–74.
- [15] Artero V, Chavarot-Kerlidou M, Fontecave M. Splitting water with cobalt. *Angew Chem Int Ed.* 2011;50(32):7238–66.
- [16] Gordon RB, Bertram M, Graedel TE. Metal stocks and sustainability. *Proc Natl Acad Sci USA.* 2006;103(5):1209–14.
- [17] Kanan MW, Surendranath Y, Nocera DG. Cobalt-phosphate oxygen-evolving compound. *Chem Soc Rev.* 2009;38(1):109–14.
- [18] Jiao F, Frei H. Nanostructured cobalt and manganese oxide clusters as efficient water oxidation catalysts. *Energy Environ Sci.* 2010;3(8):1018–27.
- [19] Wiechen M, Berends HM, Kurz P. Water oxidation catalysed by manganese compounds: from complexes to ‘biomimetic rocks’. *Dalton Trans.* 2012;41(1):21–31.
- [20] Puntiliero F, Sartorel A, Orlandi M, La Ganga G, Serroni S, Bonchio M, et al. Photoinduced water oxidation using dendrimeric Ru(II) complexes as photosensitizers. *Coord Chem Rev.* 2011;255(21–22):2594–601.
- [21] Duan LL, Tong LP, Xu YH, Sun LC. Visible light-driven water oxidation—from molecular catalysts to photoelectrochemical cells. *Energy Environ Sci.* 2011;4(9):3296–313.
- [22] Concepcion JJ, Jurss JW, Brennaman MK, Hoertz PG, Patrocino AOT, Murakami Iha NY, et al. Making Oxygen with Ruthenium Complexes. *Acc Chem Res.* 2009;42(12):1954–65.
- [23] Horizon 2020, the Framework Programme for Research and Innovation, [cited Feb. 22, 2013]. Available from: [http://ec.europa.eu/research/horizon2020/index\\_en.cfm?pg=h2020](http://ec.europa.eu/research/horizon2020/index_en.cfm?pg=h2020)
- [24] Lubitz W, Reijerse EJ, Messinger J. Solar water-splitting into H<sub>2</sub> and O<sub>2</sub>: design principles of photosystem II and hydrogenases. *Energy Environ Sci.* 2008;1(1):15–31.
- [25] Magnuson A, Anderlund M, Johansson O, Lindblad P, Lomoth R, Polivka T, et al. Biomimetic and Microbial Approaches to Solar Fuel Generation. *Acc Chem Res.* 2009;42(12):1899–909.

- [26] Herrero C, Quaranta A, Leibl W, Rutherford AW, Aukauloo A. Artificial photosynthetic systems. Using light and water to provide electrons and protons for the synthesis of a fuel. *Energy Environ Sci.* 2011;4(7):2353–65.
- [27] Cady CW, Crabtree RH, Brudvig GW. Functional models for the oxygen-evolving complex of photosystem II. *Coord Chem Rev.* 2008;252(3–4):444–55.
- [28] Dau H, Zaharieva I. Principles, Efficiency, and Blueprint Character of Solar-Energy Conversion in Photosynthetic Water Oxidation. *Acc Chem Res.* 2009;42(12):1861–70.
- [29] Rigsby ML, Mandal S, Nam W, Spencer LC, Llobet A, Stahl SS. Cobalt analogs of Ru-based water oxidation catalysts: overcoming thermodynamic instability and kinetic lability to achieve electrocatalytic O<sub>2</sub> evolution. *Chem Sci.* 2012;3(10):3058–62.
- [30] Andreiadis ES, Chavaro Kerlidou M, Fontecave M, Artero V. Artificial Photosynthesis: From Molecular Catalysts for Light-driven Water Splitting to Photoelectrochemical Cells. *Photochem Photobiol.* 2011;87(5):946–64.
- [31] Dempsey JL, Brunschwig BS, Winkler JR, Gray HB. Hydrogen Evolution Catalyzed by Cobaloximes. *Acc Chem Res.* 2009;42(12):1995–2004.
- [32] Walter MG, Warren EL, McKone JR, Boettcher SW, Mi Q, Santori EA, et al. *Chem Rev.* 2010;110:6446–73.
- [33] Maeda K, Domen K. Surface Nanostructures in Photocatalysts for Visible-Light-Driven Water Splitting. *Photocatalysis.* 2011;30:95–119.
- [34] Ellmer K, Fiechter S. Solar energetics – From photovoltaics to water splitting – Preface. *Physica Status Solidi B-Basic Solid State Physics.* 2008;245(9):1741–2.
- [35] Tran PD, Wong LH, Barber J, Loo JSC. Recent advances in hybrid photocatalysts for solar fuel production. *Energy Environ Sci.* 2012;5(3):5902–18.
- [36] Bora DK, Braun A, Constable EC. “In rust we trust”. Hematite – the prospective inorganic backbone for artificial photosynthesis. *Energy Environ Sci.* 2013.
- [37] Badura A, Kothe T, Schuhmann W, Rogner M. Wiring photosynthetic enzymes to electrodes. *Energy Environ Sci.* 2011;4(9):3263–74.
- [38] Lubitz W, Reijerse E, van Gastel M. [NiFe] and [FeFe] hydrogenases studied by advanced magnetic resonance techniques. *Chem Rev.* 2007;107(10):4331–65.
- [39] Gust D, Moore TA, Moore AL. Solar Fuels via Artificial Photosynthesis. *Acc Chem Res.* 2009;42(12):1890–8.
- [40] Hammarstrom L, Styring S. Proton-coupled electron transfer of tyrosines in Photosystem II and model systems for artificial photosynthesis: the role of a redox-active link between catalyst and photosensitizer. *Energy Environ Sci.* 2011;4(7):2379–88.
- [41] Dempsey JL, Winkler JR, Gray HB. Proton-Coupled Electron Flow in Protein Redox Machines. *Chem Rev.* 2010;110(12):7024–39.
- [42] Alia A, Buda F, de Groot H, Matysik J. Solid-State NMR of Nanomachines Involved in Photosynthetic Energy Conversion. *Annual Review of Biophysics.* 2013;42(1).
- [43] Nocera DG. The Artificial Leaf. *Acc Chem Res.* 2012;45(5):767–76.
- [44] See e.g. Building a new industrial revolution in Europe, [updated Dec. 18, 2012, cited Jan. 8, 2013]. Available from: [http://ec.europa.eu/enterprise/magazine/articles/industrial-policy/article\\_11038\\_en.htm](http://ec.europa.eu/enterprise/magazine/articles/industrial-policy/article_11038_en.htm)

## The Authors



Anders Thapper (b. 1972) did his PhD in Inorganic Chemistry at Lund University, Sweden in 2001. After a postdoctoral stay at Universidade Nova de Lisboa, Portugal, he went to Uppsala University in 2005. In Uppsala he initially studied Photosystem II together with Pr. Stenbjörn

Styring and is now a researcher working with artificial photosynthesis and manganese and cobalt based catalysts for water oxidation. He has published over 25 papers on artificial photosynthesis and bioinorganic chemistry.



Stenbjörn Styring (b. 1951) took his PhD in 1985 in biochemistry at Göteborg University and then carried out post doctoral studies in Gif-sur-Yvette and at CEA Saclay. He was chair in Biochemistry at Lund University before he moved to Uppsala University where he is a Professor in Molecular Biomimetics since 2006. His research concerns mechanistic aspects of the oxygen evolving enzyme in Photosystem II, as well as artificial photosynthesis of manganese, cobalt and ruthenium-manganese complexes intended for photocatalytic oxidation of water and he has ca 250 publications in those fields. He leads the Swedish Consortium for Artificial Photosynthesis since the start in 1994. For the EU he has coordinated the networks Ru-Mn for Artificial Photosynthesis, SOLAR-H and SOLAR-H2.




Guido Saracco (b. 1965) got his PhD in Chemical Engineering in 1995 at the Politecnico di Torino. He is now the Head of the Department of Applied Science and Technology and Chair of “Chemistry”. He authored more than 200 publications on environmental catalysis, clean energy production processes, treatment of industrial effluents, hydrogen and fuel cell technologies, biofuels, and water photolysis, a field where he coordinates the VII FP EU projects Solhydromics, ArtipHyction and Eco2CO2 . . . .



A. William Rutherford (b. 1955) got his PhD at University College London. He did post-doctoral studies in biophysics of photosynthesis at the University of Illinois USA, RIKEN Japan and CEA Saclay France. He entered the CNRS in 1983 working on water splitting in association with

the chemistry group at University Paris XI. He became head of the CNRS unit in 1992 and becoming Head of CEA Service of Bioenergetics in 2000. From 2011 he took up the Chair of Biochemistry of Solar Energy at Imperial College London, joining the Imperial's Artificial Leaf Program. He is President of the International Society of Photosynthesis Research, an EMBO member and holds a Wolfson Merit Award. His main fields are natural and artificial photosynthesis having ca. 180 papers in refereed journals.



Bruno Robert (b. 1957) graduated in theoretical physics at the University Pierre et Marie Curie (Paris 6). After a PhD in Natural Sciences and a post-doctoral stay at the Department of Chemistry of Harvard University, he developed a team on photosynthetic light-harvesting and primary charge separation processes at the CEA Saclay, France. Presently President of the French Biophysical Society, Professor of Physics at the Vrije Universiteit Amsterdam, he was in 2011 laureate of an ERC advanced grant in physics. He is in charge, together with Pr. Sebastian Fiechter, of the coordination of AMPEA Artificial Photosynthesis.



Ann Magnusson (b. 1964) took her PhD in Biochemistry at Lund University and carried out post-doctoral research at University of California at Davis and Michigan State University. Since 2005 she is an associate professor at Uppsala University. A member of the Swedish Consortium for Artificial

Photosynthesis since 1996, her research interests include manganese- and ruthenium-manganese biomimetic complexes for artificial photosynthesis, as well as the photosynthetic enzymes of plants and cyanobacteria.



Wolfgang Lubitz (b. 1949) received his doctoral degree (1977) and habilitation (1982) in Chemistry at the FU Berlin. From 1983–89 he worked at UC San Diego and as assistant and associate professor at FU Berlin, and 1989–91 at the University Stuttgart. From 1991–2001 he was Chair of Physical Chemistry at the

TU Berlin. In 2000 he became Scientific Member of the Max Planck Society and is now Director at the new Max Planck Institute for Chemical Energy Conversion in Mülheim/Ruhr. His work focuses on structure and function of energy-converting biological systems and related chemical models using spectroscopic, electrochemical and theoretical methods. The main subjects are metallo-enzymes (wateroxidase and hydrogenase). His work is documented in over 350 publications.



Antoni Llobet (b. 1960) received his BS and PhD degrees in Chemistry from the Universitat Autònoma of Barcelona in Spain. Following that, he was a postdoctoral fellow at the University of North Carolina, in the laboratories of Thomas J. Meyer and at Texas A&M University with Donald T. Sawyer

and Arthur E. Martell. He is currently a professor of Chemistry at Universitat Autònoma of Barcelona in Spain, visiting professor at Ewha Womens University in Seoul, Korea and group leader at ICIQ in Spain. His research interests are related to all aspects of redox catalysis and artificial photosynthesis.



Philipp Kurz (b. 1976) studied chemistry in Leipzig (Germany) and Zurich (Switzerland). His interest in photosynthesis started already as a diploma student, when he investigated photosynthetic picoplankton from Lake Lucerne in Switzerland. Since then, the synthesis and investigation of inorganic compounds for artificial photosynthesis has been the central topic of his research, both as a PhD student of Roger Alberto at the University of Zurich

and as a postdoc in the group of Stenbjörn Styring in Uppsala (Sweden). From 2007 to 2012 he was a junior research group leader in Kiel (Germany) and recently moved to Freiburg (Germany) where he is a professor at the Institute for Inorganic and Analytical Chemistry.



Alfred Holzwarth (b. 1949) studied chemistry at Freie Universität Berlin and at the Swiss Federal Institute of Technology, Zürich. After his PhD in Physical Chemistry, he moved to the Max-Planck-Institute for Chemical Energy Conversion in Mülheim/Ruhr, Germany to study photo-induced energy and

electron transfer reactions. He is also teaching Biophysics at the Heinrich-Heine-University in Düsseldorf. He is directing one of the leading groups in the study of primary processes and regulation mechanisms in natural photosynthesis. During the last 10 years his research focused on the transfer of knowledge from natural photosynthesis to the design of artificial photosynthetic systems. He has also been very active in science policy actions to pave the way for research activities aiming at the long-term replacement of fossil fuels by renewable fuels.



Sebastian Fiechter (b. 1954) studied mineralogy and crystallography at the Albert-Ludwigs-Universität Freiburg (Germany) and got his PhD on a topic of crystal growth supervised by Pr. R. Nitsche. In 1983 he joined the group of Pr. H. Tributsch at the Hahn-Meitner-Institute (HMI) in Berlin working

on the development of new absorber materials for thin film solar cells. From 1998–2008 he focused his research on novel fuel cell catalysts for the oxygen reduction reaction. In 2008 he introduced together with Pr. H.-J. Lewerenz (now CALTECH – JCAP; Pasadena) the research topic Solar Fuels at the Helmholtz Zentrum Berlin, successor of HMI, which has led to the foundation of a new institute directed by Pr. Dr. Roel van de Krol. Pr. Fiechter is teaching at the Technical University Berlin. Together with Pr. B. Robert he is coordinating AMPEA Artificial Photosynthesis.



Huub de Groot (b. 1958) holds a PhD in physics from Leiden University. After a period at MIT he joined the chemistry faculty in Leiden. He works with Solid State NMR on photosynthesis, on biomimetic catalysts and on multiscale modelling for the design of modular nanodevices

for solar fuel. He coordinated the ESF science policy brief on solar fuels, is in the board of the EuroSolarFuel Eurocores program and coordinates one of its collaborative research programs. Huub de Groot serves as the scientific director of the Dutch BioSolar Cells public private partnership, and contributes to its research, valorization and innovation projects.



Sebastiano Campagna (b. 1959) received the Laurea in Chemistry (cum laude) from the University of Messina in 1983. Since 1985 he spent about a decade as a post-doctoral fellow at the Chemistry Department “G. Ciamician” of the University of Bologna, in the group of Pr. Vincenzo Balzani. In 1998 he

joined the Science Faculty of the University of Messina, where he is now Professor of Physical Chemistry. In 1995 he was awarded the Raffaello Nasini Prize (Società Chimica Italiana). He is the Coordinator of the Nano-Solar network project, funded by the Italian MIUR, and co-Director of SOLAR-CHEM, an Italian interuniversity center for artificial photosynthesis. His research fields include light- and redox-active dendrimers, Ru(II) photophysics, and artificial photosynthesis. He is the author of ca. 200 papers.



Artur Braun (b. 1965) is a physicist from RWTH Aachen (magnetic surfaces) with a doctoral degree in electrochemistry from ETH Zürich and Paul Scherrer Institut (supercapacitors). Artur worked with Pr. Elton Cairns and Pr. Stephen P. Cramer at Lawrence Berkeley National Laboratory

(lithium batteries and protein spectroscopy) and with Pr. Gerald P. Huffman at University of Kentucky (fossil fuels), before he returned to Empa in Switzerland as Marie Curie

Fellow and group leader (fuel cells, solar hydrogen). He has a prestigiously low Erdős number of 3 and authored over 100 publications. Due to his expertise in electronic structure and transport properties of energy materials and synchrotron/neutron methods, he is energy editor of Current Applied Physics (Elsevier). Artur Braun considers artificial photosynthesis a chance, not a challenge.



Hervé Bercegol (b. 1966) is a graduate of the École Normale Supérieure (Paris). He got a doctorate in physics from the University Pierre et Marie Curie (Paris 6) in 1991 for an experimental, fundamental work on a 2D solid. After a postdoctoral work on surface

spectroscopies of materials in the University of Cincinnati USA, he worked for the company Saint-Gobain and for CEA, mainly on surface related materials questions. From 1998 to 2008, he was the leader of laser damage studies for the Laser Mégajoule project. Since 2010, he has been in charge of a new CEA programme designed to foster innovative concepts for

energy efficiency and renewable energy. Within the European Energy Research Alliance, Hervé Bercegol coordinates the Joint Programme AMPEA.



Vincent Artero (b. 1973) is a graduate of the École Normale Supérieure (Paris) and from the University Pierre et Marie Curie (Paris 6). His doctoral work (2000) under the supervision of Prof. A. Proust dealt with organometallic derivatives of polyoxometalates. After a postdoctoral stay in Aachen with Prof. U. Kölle, he joined in 2001 the Laboratory of Chemistry and Biology of Metals in Grenoble where he obtained a position in the Life Science Division of the CEA. His current research interests are in the structural and functional modelisation of hydrogenases and the design of artificial systems for the photo- and electroproduction of hydrogen. He obtained in 2011 the Grand Prix Mergier-Bourdeix from the French Academy of Sciences and is a 2012 laureate of the ERC starting grant fellowship.

